

Increasing safety for youth in and leaving foster care

Report to Blue Ribbon Commission on Child
Protection – February 21, 2014

Wendy B. Smith, Ph.D., LCSW

Associate Dean, Faculty Development &

Associate Clinical Professor

Topics

- Introduction
- Impact of maltreatment
- Use of psychotropic medications
- Suicidality among youth
- Commercially sexually exploited children (CSEC) and foster care
- Recommendations

Impact of maltreatment

- 40%-60% of maltreated children have mental health problems (Linares et al, 2013).
- Self-regulation
 - Executive functioning
 - Managing strong feelings and impulses
- Social-emotional processing
 - Compromised interpersonal relationships

Dysregulated stress response system

- Trained at biological level to be on alert
- Can't turn off trauma-induced activation of stress response system
- Once adaptive and protective
- Now maladaptive and can increase dangers

Impact on development over time

- Influence on brain, social, and identity development going forward
- Compounded by broken or disturbed attachments
- Unresolved trauma and unresolved grief add to internal burden, creating risks from others and from the self.

Poor mental health presents risks to safety

- Contributes to:
 - to school failure
 - Unemployment
 - Poverty
 - Early parenthood
 - Lack of medical treatment
 - Homelessness
 - Victimization and exploitation
 - Suicide
 - Violence to others

Psychotropic medication and children in foster care

- Use of mental health services 15-20 times that of children in general population (Wolff et al, 2013)
- 40%-60% meet criteria for at least one psychiatric disorder (Wolff et al, 2013)
- By adolescence, 63% have at least one diagnosis; 23% have three or more. (White et al, 2007)
- Psychotropic meds: 13%-40%, compared with 4%-6% in general population. (Wolff et al, 2013; Lyons et al, 2013; Burcu et al, 2013)

- Polypharmacy (multiple medications concurrently) – children in care take greater average number of medications (Lyons et al, 2013)
- More frequently prescribed antipsychotics, especially for ADHD, than general population. (Linares, 2013)
- Nearly 9 times rate of Medicaid-enrolled children not in care (USDHHS, 2012)

Which children?

- Those in most restrictive placement settings
- White males
- Males with behavior problems
- Use and use of multiples higher among 6- to 10-year-olds than 11- to 17-year-olds
- Those **under 1 year** have 0.3%-2/1%, compared with 0.1%-1.2% of those not in **care**. (USDHHS, 2012)

Safety risks in adolescence: suicidality

- Youth in child welfare and juvenile justice systems 3-5 times more likely to die by suicide (National Center for Prevention of Youth Suicide, 2013).
- Children and youth in care at greater risk of suicide and attempting suicide (Katz et al, 2011)
- Trauma involving assaultive violence and child sexual abuse predicts suicidal behavior.
- Depressive symptoms play major role.

Suicidality: which youth?

- Deaths: males
 - 10-14 yrs: 3.8 to 1
 - 15-19 yrs: 4.7 to 1
 - 20-24 yrs: 6.7 to 1.
- Attempts: females more frequent in teen years
- Those with foster care histories 4 times more likely to report suicidal ideation.
- Homelessness and child sexual abuse increase risk (Corbett et al, 2012)
- Among maltreated 8-yr-olds: 10% reported wanting to kill themselves.

Safety risks in adolescence: sex trafficking

- 50% of those sold into trafficking in CA and 85% in NY have child welfare
- Among survivors of trafficking, 98% had cw involvement
- High rates of sexual assault in childhood.
- Recruited at younger ages
- Male, female, ethnically diverse, LGBT (Lillie, 2013; Spangenberg, 2001)

Sex Trafficking and Homelessness

- 22% of youth aging out become homeless
- 75% of all victims were at one point homeless
- 1 in 3 homeless teens lured into CSE within 48 hours.
- Recruited youth have aged out, run away, or been lured away from home.

Recommendation #1

- **Use of psychotropic medications should be carefully considered and based on screening, assessment, and treatment planning.**
 - Consider availability of nonpharmacological interventions
 - Weigh risks and benefits of psychotropics in children
 - Use should always entail effective medication monitoring, and informed, shared decision-making and ongoing communication.
 - Mental health experts, including board certified psychiatrists, should provide consultation when establishing processes of consent and monitoring.

Recommendation #2

- **Screenings** (for depression, for trauma and status of trauma resolution, for cognitive, behavioral, emotional, and social development relative to normative development) should take place **prior to entry into care, at entry, at regular intervals, during transition planning, and at exit from care.**

Recommendation #3

- **When psychotherapy is indicated, tailor treatment to individual youth rather than disorder.**
 - Consider symptoms, strengths, vulnerabilities, culture, history, developmental status, current situation/environment, available treatments, evidence of effectiveness.
 - Consider gender, ethnicity, and culture
 - Address trauma- and attachment-related issues.

Recommendation #4

- Ensure the **maximum placement stability** and, with cooperation from school districts, **school stability, especially during middle school (a time of high sensitivity to changes in environment)** and high school.

Recommendation #5

- Begin **planning for post-secondary education or career training** during middle school.
 - Monitor educational progress toward goals at regular intervals.
- **Academic achievement is a protective factor**, building self-esteem, positive identity, and future employment potential.
 - Attendance at 4-year college prevents homelessness.

Recommendation #6

- **Sex trafficking prevention:**
 - **Document** nature and extent of problem in LA County
 - **Increase awareness** of problem
 - **Train law enforcement and social workers** to recognize possible indicators.
 - **Involve youth as partners** in developing policies, programs, and resources.

Recommendation #7

- **Transition planning with youth:**
 - Careful assessment of availability of **caring others and social support network, including cultural and spiritual dimensions**
 - Remediation of gaps prior to exit.
 - **Network assessment should include biological and extended family** (potential sources of strength, but often continuing stressors and risks to fragile well-being.
 - Provide services aimed at ameliorating family relationships prior to exit from care.

Recommendation # 8

- **Build in safeguards against homelessness** following discharge from care
 - Onsite visits to planned living arrangement.
 - No discharge without certainty of housing.
 - Monitoring of housing stability for the year (or more) following discharge.

References

- .
- Ai, A., Foster, L., Pecora, P., Delaney, N. & Rodriguez, W. (2013) Reshaping child welfare's response to trauma: Assessment, evidence-based intervention, and new research perspectives. *Research on Social Work Practice*, 23(6), 65.
-
- Anderson, H. (2011). Suicide ideation, depressive symptoms, and out-of-home placement among youth in the U.S. child welfare system. *Journal of Clinical Child & Adolescent Psychology*, 40(6), 790.
- Burcu, M. Zito, J., Safer, D. & Ibe, A. (2013) Psychotropic medication patterns in medicaid-insured youth based on clinician-reported maltreatment status . *J of Child and Family Studies*, (1062-1024).
- Chavira, D., Accurso, E., Garland, A. & Hough, R. (2010). Suicidal behavior among youth in five public sectors of care. *Child and Adolescent Mental Health*, 15(1), 44.

References, cont'd.

-
- Corbett, R., Mazin, N., Grimshaw, R. & Bebbington, P. (2012) Thinking of suicide: understanding the risks associated with child institutional care. *Criminal Justice Matters*, 90(1), 38.
- Fawley-King, K. & Merz, E. (2014). Effects of child maltreatment on brain development. In Matto, H., Strolin-Goltzman, J. & Ballan, M., Eds. *Neuroscience for social work: Current research and practice*. New York: Springer Publishing Company.
- Katz, L., Au, W., Singal, D. et al (2011). Suicide and suicide attempts in children and adolescents in the child welfare system. 183(17), 1977.
- Lillie, M. (2013). An unholy alliance: The connection between foster care and human trafficking. Fifth Annual Interdisciplinary Conference on Human Trafficking The OLP Foundation and Human Trafficking Search. *Net*. Retrieved 2-5-14 from <http://digitalcommons.unl.edu>
-

References, cont'd.

- Linares, L., Martinez-Martin, N. & Castellanos, F. (2013). Stimulant and atypical antipsychotic medications for children placed in foster homes. *PLoS ONE*, 8(1), e54152.
-
- Lyons, C., Wasserman, G., Olsson, M., McReynolds, L., Musabegovic, H. & Keating, J. (2013) Psychotropic medication patterns among youth in juvenile justice. *Administration and Policy in Mental Health*, 40, 58-68.
-
- Narendorf, Sarah. (2013) Psychotropic medication use patterns in older foster youth. *Presentation at 13th Annual Conference of Society for Social Work and Research*. San Diego, CA.
- National Center for the Prevention of Youth Suicide. (2013) Preventing suicidal behavior among youth in foster care. Retrieved 2-5-14 from http://www.suicidology.org/c/document_library/get_file?folderId=261&name=DLFE-557.pdf
-

References, cont'd.

- Smith, W. (2011). Youth leaving foster care: A developmental, relationship-based approach to practice. New York: Oxford University Press.
-
- Smith, W. (2014). The role of neurobiology in social work practice with youth transitioning from foster care. In Matto, H., Strolin-Goltzman, J. & Ballan, M., Eds. *Neuroscience for social work: Current research and practice*. New York: Springer Publishing Company.
-
- Spangenberg, M. (2001). Prostituted youth in New York City: An overview. ECPAT-USA. Retrieved 2-17-14 from http://www.resources.ecpat.net/EI/EI_publications.asp
- Taliaferro, L. & Muehlenkamp, J. (2013) Risk and protective factors that distinguish adolescents who attempt suicide from those who only consider suicide in the past year. *Suicide and Life-Threatening Behavior*. Doi: 10.1111/sltb.12046.
-

References, cont'd.

- U.S. Department of Health and Human Services, Administration on Children, Youth, and Families (USDHHS) a. (2012) Information Memorandum, Log No ACYF-CB-IM-12-03. Retrieved 2-16-14 from <http://www.acf.hhs.gov/sites/default/files/cb/im1203.pdf>
-
- U.S. Department of Health and Human Services, Administration on Children, Youth, and Families (USDHHS) b. (2012) Information Memorandum, Log No ACYF-CB-IM-12-04. Retrieved 2-10-14 from <http://www.acf.hhs.gov/sites/default/files/cb/im1204.pdf>
-
- White, CR, Havalchack, A., Jackson, L., O'Brien, K., & Pecora, P. (2007) Mental health, ethnicity, sexuality, and spirituality among youth in foster care: Findings from the Casey Field Office Mental Health Study. Seattle, WA: Casey Family Programs.
-
- Wolff, J., Carleton, R. & Drilea, S. (2013) Are rates of psychotropic medication use really higher among children in child welfare? *26th Annual Children's Mental Health Research and Policy Conference*. Tampla, Fla.
-